



Targeted sequencing and integrative analysis to prioritize candidate genes in neurodevelopmental disorders

Yi Zhang¹ · Tao Wang^{2,3,4} · Yan Wang³ · Kun Xia^{2,5,6} · Jinchen Li^{1,2,7} · Zhongsheng Sun^{3,8,9,10} 

Received: 15 December 2020 / Accepted: 29 March 2021
© The Author(s) 2021

Abstract

Neurodevelopmental disorders (NDDs) are a group of diseases characterized by high heterogeneity and frequently co-occurring symptoms. The mutational spectrum in patients with NDDs is largely incomplete. Here, we sequenced 547 genes from 1102 patients with NDDs and validated 1271 potential functional variants, including 108 de novo variants (DNVs) in 78 autosomal genes and seven inherited hemizygous variants in six X chromosomal genes. Notably, 36 of these 78 genes are the first to be reported in Chinese patients with NDDs. By integrating our genetic data with public data, we prioritized 212 NDD candidate genes with FDR < 0.1, including 17 novel genes. The novel candidate genes interacted or were co-expressed with known candidate genes, forming a functional network involved in known pathways. We highlighted *MSL2*, which carried two de novo protein-truncating variants (p.L192Vfs*3 and p.S486Ifs*11) and was frequently connected with known candidate genes. This study provides the mutational spectrum of NDDs in China and prioritizes 212 NDD candidate genes for further functional validation and genetic counseling.

Keywords Neurodevelopmental disorders · Targeted sequencing · De novo variants · Candidate genes

Introduction

Neurodevelopmental disorders (NDDs) are complex and heterogeneous disorders characterized by impaired motor function, learning, and verbal and non-verbal communication resulting from dysfunction during brain development [1–4]. NDDs span a very wide range of neurological and psychiatric

disorders [1, 5] and affect more than 3% of children worldwide [6]. Although each diagnosis is distinct in clinical settings, in general, NDDs are characterized by high clinical and genetic heterogeneity. Previous studies have estimated the heritability of NDDs, with the highest being up to ~ 80% [7–10]. Multiple phenotype-genotype correlation studies have revealed that patients carrying deleterious variants in the same

Yi Zhang and Tao Wang contributed equally to this work.

✉ Kun Xia
xiakun@sklmg.edu.cn

✉ Jinchen Li
lijinchen@csu.edu.cn

✉ Zhongsheng Sun
sunzs@biols.ac.cn

¹ National Clinical Research Centre for Geriatric Disorders, Department of Geriatrics, Xiangya Hospital, Central South University, Changsha 410083, Hunan, China

² Center for Medical Genetics and Hunan Key Laboratory of Medical Genetics, School of Life Sciences, Central South University, Changsha 410083, Hunan, China

³ Beijing Institutes of Life Science, Chinese Academy of Sciences, Beijing 100101, China

⁴ DIAGenes Precision Medicine, Beijing 102600, China

⁵ CAS Center for Excellence in Brain Science and Intelligent Technology (CEBSIT), Shanghai 200031, China

⁶ School of Basic Medical Science, Central South University, Changsha 410078, Hunan, China

⁷ Department of Neurology, Xiangya Hospital, Central South University, Changsha 410008, Hunan, China

⁸ Institute of Genomic Medicine, Wenzhou Medical University, Wenzhou 325035, China

⁹ CAS Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing 100049, China

¹⁰ State Key Laboratory of Integrated Management of Pest Insects and Rodents, Chinese Academy of Sciences, Beijing 100101, China

risk gene have manifested broad clinical phenotypes, including impaired social interaction, developmental delay, regression, and repetitive behavior [11–13]. The high level of heterogeneity and similar pathogenetic mechanisms present challenges for clinical diagnosis and treatment.

Recent studies have reported that targeted sequencing is a powerful and cost-effective tool for discovering new genetic risk genes in many diseases, particularly diseases with high genetic heterogeneity [4, 14, 15]. Both rare inherited and de novo variants (DNVs) have been demonstrated to contribute to NDDs [16–18]. Stessman et al. sequenced 208 candidate genes in patients with NDDs and identified 91 risk genes with locus-specific significance for disruptive variants in 5.7% of patients [4]. Wang et al. performed targeted sequencing of 189 autism spectrum disorder (ASD) candidate risk genes in Chinese patients and found that genes identified in European ASD cohorts were highly relevant in Chinese cohorts [19]. Guo et al. expanded the above study to a larger cohort of patients and provided support for a multifactorial model of ASD risk [20]. Some studies have shown that there are multiple risk genes that are shared between neurodevelopmental disorders, and integrating datasets at multiple levels is beneficial for the etiology of NDDs [21, 22]. Takata et al. combined published DNV data and found that integrative analyses was conducive to identifying significant genes and extending ASD-related molecular and brain networks [23]. Gonzalez-Mantilla et al. used multilevel data-integration approach and identified novel candidate genes for developmental brain disorders [24]. We have previously demonstrated that DNVs, gene set enrichment analysis, and protein-protein interaction (PPI)/co-expression analysis can provide new insights into the genetic mechanisms underpinning NDDs [25–28]. However, only a few known pathogenetic genes have been described to explain the genetic causes of NDDs, and the etiology behind NDDs remains unclear.

Here, we used targeted sequencing to examine 547 genes from 1102 Chinese patients with NDDs. We identified potential functional variants and explored the patterns of DNVs in Chinese patients with NDDs. We then integrated our dataset with public datasets of neurodevelopmental disorders from the Gene4Denovo [29] database to prioritize NDD candidate genes and discover novel candidate genes. Finally, we investigated whether novel candidate genes were functionally associated with known candidate genes using multilevel bioinformatics analysis.

Materials and Methods

Panel Gene Design and Targeted Sequencing

In this study, potential NDD risk genes are collected based on the following criteria: (1) genes recorded in the Online

Mendelian Inheritance in Man (OMIM) (<https://omim.org/>) are associated with NDDs, mainly including autism, cerebral palsy, mental retardation, and epilepsy; (2) candidate genes in the NPdenovo database [28]; (3) strong risk genes from the SFARI Gene database (<https://gene.sfari.org/>) [30]; and (4) genes prioritized in our previous NDD-related studies [25, 26, 31, 32]. After removing duplicated genes, 547 target genes were selected (Table S1).

A total of 935 unrelated trios (proband and their unaffected parents) and 167 probands without parents were recruited from China. Genomic DNA (1 µg) extracted from whole blood was sheared and assembled into a DNA library prior to targeted sequencing. The Illumina X10 sequencing system (Illumina, San Diego, CA, USA) was used to generate paired-end raw data. This panel resulted in an average depth of 181.46× in target regions, and 98.82% of target bases were covered with depth $\geq 10\times$ on average. This study was approved by the Institutional Review Board of the State Key Laboratory of Medical Genetics, School of Life Sciences, Central South University, Changsha, Hunan, China. All subjects who participated in this study provided informed consent prior to sample collection.

Variation Detection and Annotation

Quality control of the sequencing data was performed using Cutadapt [33] and FastQC (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) to remove adapter and unqualified sequences, respectively. BWA-MEM [34] was employed to align the clean reads to the human reference genome (hg19). Samtools [35] utilities were used to mark duplicate reads and generate position-sorted files, while the Genome Analysis Toolkit HaplotypeCaller was used to call variants. Comprehensive annotation of all variants was performed using ANNOVAR [36], including functional implications (gene region, functional effect, mRNA GeneBank accession number, amino acid change, cytoband, etc.), functional predictions for missense variants, and allele frequencies of gnomAD, ExAC, and in-house data (1113 WES samples and 2469 WGS samples). Deleterious missense variants (Dmis) were predicted by ReVe [37] and missense variants with ReVe ≤ 0.7 were excluded. Only protein-truncating variants (PTVs, including stop-gain, stop-loss, frameshift, and splicing) and Dmis with minor-allele frequencies $\leq 0.1\%$ were defined as potential functional variants and selected for further analysis. Sanger sequencing was used to validate all potential functional variants in our study.

Prioritization of Candidate Genes

Using de novo PTVs, together with Dmis and background DNV rates, we employed the TADA classification tool [38]

to prioritize candidate genes, with gene exhibiting a false discovery rate (FDR) < 0.1 defined as candidates. First, DNVs from 935 trios, inherited variants from 1102 probands, and genetic variants from 3582 in-house Chinese controls (Table S6) were applied to TADA model, and we prioritized 17 candidate genes with FDR values < 0.1. To increase the statistical power, we integrated DNVs from 16,807 probands with different types of NDDs and included 3391 controls from the Gene4Denovo database (version 1.0) [39] and 208 candidate genes with FDR values < 0.1 for further analysis. All samples integrated in our study have been carefully deduplicated.

To discover novel candidate genes, we collected known candidate genes from previous studies and related databases, and excluded them from the list of candidate genes prioritized by TADA. Known candidate genes were defined as follows: (1) genes defined as risk genes in any ten recent publications involving large-scale exome, whole-genome, or targeted sequencing (De Rubeis et al. [17], Sanders et al. [40], Lelieveld et al. [41], McRae et al. [42], Stessman et al. [43], Yuen et al. [44], Nguyen et al. [22], Takata et al. [23], Coe et al. [21], and Satterstrom et al. [45]); (2) genes collected from OMIM that were associated with neurodevelopmental disorders, including intellectual developmental disorder, autism, mental retardation, epileptic encephalopathy, and schizophrenia; (3) genes belonging to the category of syndromic gene or score category of 1 or 2 in SFARI Gene [30]. Genes that met any of the above conditions were classified as known candidate genes. In contrast, genes that failed to meet any of the above conditions above and had no obvious genetic evidence in association with NDDs in PubMed were classified as novel candidate genes.

Permutation Test

Spatial and temporal expression data of the human brain were downloaded from BrainSpan (<http://www.brainspan.org/>). For ethical reasons, we removed samples during fetal stages and selected postnatal cortical samples for further study. We calculated the Pearson correlation coefficients between any two genes based on their expression levels. Gene pairs with $|R| > 0.7$ were regarded as being co-expressed in the human brain. The permutation test was performed to compare the novel and the known candidate gene sets, in order to evaluate their functional connections. In brief, we compared the number of co-expressed genes within the novel and the known candidate gene sets and their connections with a random gene set of 1,000,000 random iterations.

Functional Network Analysis

We downloaded PPI data from IntAct Molecular Interaction database (<https://www.ebi.ac.uk/intact/>) for analysis and

considered gene pairs with an intact-miscore ≥ 0.45 to be interacting genes. We then constructed a network based on gene pairs selected from our PPI and co-expression analyses. Novel candidate genes were defined as “seed genes” that were directly connected and used to form an interconnected functional network. Known NDD candidate genes directly connected to at least two novel candidate genes were added to the above network. To further investigate the functional pathways of novel candidate genes, we performed GO enrichment analysis using MetaScape (<https://metascape.org>). Similar pathways were merged into a single cluster. Network figures were drawn using Cytoscape v.3.7.2.

Results

DNVs in Our Chinese Cohort

In this study, we sequenced 547 target genes in 1102 Chinese patients with NDDs and identified a set of predicted potential functional variants, including PTVs and Dmis (Fig. 1 and Table S1). Using Sanger sequencing, we successfully validated 1271 potential functional variants, including 108 de novo variants (54 de novo PTVs and 54 de novo Dmis), 975 inherited variants (156 inherited Dmis and 819 inherited PTVs), and 188 undetermined variants (36 PTVs and 152 Dmis) (Fig. S1 and Table S2). We found that 108 DNVs in 78 genes appeared in approximately 10.91% (102/935) of patients in our Chinese cohort. Among these 78 genes, 36 genes with DNVs are the first to be reported in Chinese patients. Furthermore, we revealed that 21 genes carried multiple DNVs, including 28 de novo PTVs and 23 de novo Dmis (Table 1). Both *SCN2A* and *MECP2* were the most frequently mutated genes, each carrying one de novo PTV and three de novo Dmis (Table 1). Five genes (*MED13L*, *GRIN2B*, *KCNQ2*, *CTNNA1*, and *TCF20*) carried three DNVs in our Chinese cohort, while another 14 genes (*ASH1L*, *SATB2*, *NRXN1*, *BCL11A*, *ADNP*, *SHANK3*, *MSL2*, *SYNE1*, *SYNGAP1*, *BRAF*, *GATAD2B*, *LLGL1*, *SLC2A1*, and *KDM5C*) carried two DNVs (Table 1). We manually collected 805 known candidate genes that have either been reported in studies on large NDD cohorts or have strong evidence associating them with NDDs in the OMIM or SFARI Gene databases [17, 21–23, 40–45] (Table S3). We found that of 21 genes with multiple DNVs that we identified, 20 are classified as known candidate genes (Table S4). For example, *TCF20*, which carries three *de novo* PTVs (p.S1803Vfs*6, p.C1795Wfs*13, and p.R1907X), was first reported in a Chinese cohort. Interestingly, we identified a potential novel NDD risk gene (*MSL2*) that carries two *de novo* PTVs

Table 1 Genes with multiple DNVs in our probands

Gene symbol	DNVs (<i>n</i> = 935)	Inherited or state unknown (<i>n</i> = 1102)	DNVs from Gene4Denovo (<i>n</i> = 16,807)	RVIS (percentile)	pLI (percentile)	Gene function
<i>SCN2A</i>	1 PTV, 3 Dmis	1 PTV, 3 Dmis	18 PTVs, 32 Dmis	- 2.51 (1.05%)	1.00 (0.72%)	Nervous system development
<i>MECP2</i>	1 PTV, 3 Dmis	1 Dmis	9 PTVs, 10 Dmis	- 0.32 (30.25%)	0.70 (24.62%)	Chromatin binding
<i>MED13L</i>	3 PTVs	2 Dmis	18 PTVs, 11 Dmis	- 1.61 (3.34%)	1.00 (0.57%)	Brain development
<i>GRIN2B</i>	1 PTV, 2 Dmis	2 Dmis	4 PTVs, 15 Dmis	- 2.48 (1.10%)	1.00 (2.19%)	Nervous system development
<i>KCNQ2</i>	1 PTV, 2 Dmis	1 PTV, 3 Dmis	2 PTVs, 25 Dmis	- 1.25 (5.97%)	1.00 (5.39%)	Regulation of neuronal excitability
<i>CTNNB1</i>	3 PTVs	1 Dmis	19 PTVs	- 1.07 (8.10%)	1.00 (4.20%)	Wnt signaling
<i>TCF20</i>	3 PTVs	3 Dmis	11 PTVs	- 3.82 (0.34%)	1.00 (2.84%)	Wnt signaling
<i>ASH1L</i>	2 PTVs	3 Dmis	7 PTVs	- 3.89 (0.31%)	1.00 (0.16%)	Chromatin organization
<i>SATB2</i>	2 PTVs	-	11 PTVs, 5 Dmis	- 1.31 (5.37%)	1.00 (4.97%)	Chromatin binding
<i>NRXN1</i>	2 Dmis	3 PTVs, 11 Dmis	2 PTVs, 5 Dmis	- 1.88 (2.30%)	1.00 (3.22%)	Nervous system development
<i>BCL11A</i>	2 PTVs	1 Dmis	6 PTVs, 3 Dmis	- 1.67 (3.07%)	0.83 (20.63%)	Brain development
<i>ADNP</i>	2 PTVs	5 Dmis	26 PTVs, 1 Dmis	- 1.54 (3.72%)	1.00 (5.98%)	Chromatin binding
<i>SHANK3</i>	2 PTVs	2 PTVs	9 PTVs	-	1.00 (4.13%)	Nervous system development
<i>MSL2</i>	2 PTVs	1 Dmis	1 PTV	- 0.72 (15.01%)	0.89 (18.19%)	Chromatin organization
<i>SYNE1</i>	2 Dmis	4 PTVs, 16 Dmis	3 Dmis	- 1.10 (7.70%)	3.75E-27 (99.45%)	Nucleotide binding
<i>SYNGAP1</i>	1 PTV, 1 Dmis	1 Dmis	31 PTVs, 1 Dmis	- 2.30 (1.36%)	1.00 (1.79%)	Postsynaptic signaling
<i>BRAF</i>	2 Dmis	1 Dmis	10 Dmis	- 0.97 (9.63%)	1.00 (2.83%)	Calcium ion binding
<i>GATAD2B</i>	2 PTVs	2 Dmis	11 PTVs	- 0.65 (16.92%)	1.00 (6.34%)	Chromatin remodeling
<i>LLGL1</i>	2 Dmis	2 Dmis, 2 PTV	1 Dmis	- 1.70 (2.94%)	0.98 (12.10%)	Axon development
<i>SLC2A1</i>	2 Dmis	1 Dmis	4 PTVs, 1 Dmis	- 0.92 (10.49%)	0.94 (15.56%)	Brain development
<i>KDM5C</i>	2 Dmis	-	3 PTVs, 2 Dmis	- 2.71 (0.88%)	1.00 (7.02%)	Chromatin organization

PTVs, protein-truncating variants, including frameshift, splicing, stop-gain, and stop-loss; Dmis, deleterious missense variants with ReVe score > 0.7


(p.S560Ifs*11 and p.L266Vfs*3). This study is the first to report an association between *MSL2* and NDDs.

We then employed probability of loss-of-function intolerance (pLI) [46] analysis from ExAC and residual variation intolerance scores (RVIS) [47] to evaluate the functional impact of genes with multiple DNVs. According to the scores of pLI and RVIS, we ranked them and used percentiles as an indicator of gene intolerance. Genes with low-percentile RVIS or pLI were more likely to be intolerant to genetic variants. As expected, it was shown that genes with multiple DNVs both exhibited significantly lower percentile pLI ($p = 4.10 \times 10^{-10}$, two-tailed Wilcoxon rank-sum test) and RVIS ($p = 1.55 \times 10^{-10}$, two-tailed Wilcoxon rank-sum test) than background genes (Fig. S2a). In particular, 20 of the 21 multiple DNV genes ranked in the top 50% of pLI and RVIS (Table 1), suggesting that they are less tolerant of damaging variants.

Inherited X-Linked Hemizygous Variants

We identified seven inherited X-linked hemizygous variants in six genes (Table S5). Among these seven variants, three variants, including two Dmis (p.R270C on *PTCHD1* and p.R197H on *SLC16A2*) and one splicing variant (c.1030-1G>C on *SLC9A7*), were recorded in the dbSNP or gnomAD database. The remaining four variants, including three Dmis (p.Q153R on *ATP6AP2*, p.R287C on *ARHGEF9*, and p.C517Y on *PLXNA3*) and one frameshift (p.N136Kfs*30 on *SLC16A2*) are reported here for the first time. Compared with the 805 known candidate genes, we note that five of the above genes are classified as known candidate genes and three (*PTCHD1*, *ATP6AP2*, and *SLC9A7*) are reported for the first time in Chinese patient with NDDs. In addition, we identified a novel candidate gene (*PLXNA3*) carrying a novel hemizygous missense

Fig. 1 Study workflow. This study consisted of four parts: (1) sample collection; (2) identification and validation of variants; (3) prioritization of candidate genes; (4) functional network analysis of novel and known candidate genes. PTVs, protein-truncating variants; Dmis, deleterious missense variants; Dmis, deleterious missense variants



variant (p.C517Y). A previous study has described an ASD patient from Maghreb carrying a rare inherited missense variant (p.D863E) in *PLXNA3* [48], suggesting that it may contribute to autism.

Prioritization of Candidate Genes

Previous studies have demonstrated that both DNVs and rare inherited variants (RIVs) contribute significantly to NDDs [49–54]. Therefore, we integrated data on DNVs and RIVs and employed TADA [38] to prioritize candidate genes in NDDs (Table S6). Based on DNVs and RIVs identified in our study, we prioritized 17 candidate genes (*SCN2A*, *KCNQ2*, *SATB2*, *SHANK3*, *GATAD2B*, *NRXN1*, *MED13L*, *SYNGAP1*, *BRAF*, *MECP2*, *GRIN2B*, *TCF20*, *CTNBN1*, *ASH1L*, *ADNP*, *BCL11A*, and *LLGL1*) with FDR values < 0.1 (Table S7). To increase the power of candidate gene detection [17, 22, 43, 55–57], we integrated data from public datasets of NDD cohorts ($N = 16,807$) with that of controls ($N = 3391$) from the Gene4Denovo [39] database (Table S8),

and prioritized 208 candidate genes with FDR values < 0.1, in which 193 candidate genes reached an FDR < 0.05 (Table S9). Together with the six genes characterized by inherited X-linked hemizygous variants, in total, we prioritized 212 NDD candidate genes, with 17 genes defined as novel candidate genes (including 13 genes with FDR values < 0.05, three genes with $0.05 \leq \text{FDR} < 0.1$ (Table 2), and one gene with an inherited X-linked hemizygous variant). We observed that the 212 candidate genes prioritized in our study exhibited significantly lower percentile pLI scores ($p < 2.20 \times 10^{-16}$, two-tailed Wilcoxon rank-sum test) and RVIS ($p < 2.20 \times 10^{-16}$, two-tailed Wilcoxon rank-sum test; Fig. S2b), consistent with the result regarding genes with multiple DNVs. A similar result was observed among the 17 novel candidate genes ($p = 0.0394$ and $p = 1.74 \times 10^{-4}$; two-tailed Wilcoxon rank-sum test; Fig. S2c), suggesting that these novel NDD candidate genes were likely to be intolerant of functional variants.

In addition, we found that DNVs and inherited X-linked hemizygous variants in 212 candidate genes were identified

Table 2 Novel candidate genes prioritized by TADA analysis

Gene symbol	DNVs (<i>n</i> = 935)	Inherited or state unknown (<i>n</i> = 1102)	DNVs from Gene4Denovo (<i>n</i> = 16,807)	FDR	RVIS (percentile)	pLI (percentile)	Gene summary
<i>SMAD6</i>	-	1 Dmis	3 PTVs, 1 Dmis	4.83E-04	-	6.97E-06 (78.89%)	Developmental and cellular process
<i>SPG7</i>	-	3 PTVs, 3 Dmis	5 Dmis	8.89E-04	- 0.91 (10.60%)	1.38E-18 (98.28%)	Anterograde axonal transport
<i>MSL2</i>	2 PTVs	1 Dmis	1 PTV	9.62E-04	- 0.72 (15.01%)	0.90 (18.20%)	Chromatin organization
<i>CYP27C1</i>	-	2 PTVs, 3 Dmis	1 PTV, 3 Dmis	1.39E-03	- 0.89 (11.06%)	1.85E-04 (70.01%)	Retinal metabolic process; retinol metabolic process
<i>ITSN1</i>	-	2 Dmis	3 PTVs, 2 Dmis	1.89E-03	- 2.89 (0.76%)	1.00 (0.87%)	Regulation of modification of postsynaptic actin cytoskeleton
<i>PSD3</i>	-	1 PTV, 2 Dmis	3 PTVs, 1 Dmis	2.11E-03	0.09 (56.40%)	0.85 (19.80%)	Nervous system development
<i>POLR3A</i>	1 Dmis	5 Dmis	2 PTVs, 1 Dmis	2.99E-03	- 2.26 (1.40%)	7.39E-14 (96.18%)	Nervous system development
<i>UBR3</i>	-	-	3 PTVs	5.41E-03	-	1.00 (5.02%)	Chromatin organization; nervous system development
<i>GALNT18</i>	-	2 Dmis	2 PTVs, 1 Dmis	7.34E-03	- 1.07 (8.11%)	4.77E-02 (47.21%)	Metabolism of proteins
<i>DCX</i>	-	-	2 PTVs, 1 Dmis	7.43E-03	- 0.67 (16.20%)	0.86 (19.34%)	Axoneme assembly
<i>LRR4</i>	-	-	1 PTV, 2 Dmis	2.56E-02	- 0.84 (11.90%)	0.88 (18.68%)	Nervous system development
<i>SPAG9</i>	1 PTV	3 Dmis	1 PTV	3.15E-02	- 1.20 (6.59%)	1.00 (1.67%)	Spinocerebellar ataxia 35
<i>ST3GAL6</i>	-	-	2 PTVs	3.23E-02	0.07 (55.10%)	1.06E-03 (64.45%)	Pre-NOTCH Expression and processing
<i>YTHDC1</i>	-	-	2 PTVs	8.48E-02	- 0.61 (18.16%)	1.00 (1.0864%)	Chromatin regulation/acetylation
<i>RRAGC</i>	-	1 Dmis	2 Dmis	9.00E-02	- 0.25 (34.05%)	0.602 (27.41%)	Regulation of TORC1 signaling
<i>DNAH17</i>	1 Dmis	3 PTVs, 16 Dmis	1 PTV, 3 Dmis	9.58E-02	-	-	Developmental process

PTVs, protein-truncating variants, including frameshift, splicing, stop-gain, and stop-loss; Dmis: deleterious missense variants with ReVe score > 0.7

in ~ 10.59% (99/935) and ~ 0.94% (7/745) of patients in our study, respectively (Table 3). DNVs and inherited X-linked hemizygous variants in known candidate genes accounted for ~ 10.05% (94/935) and ~ 0.81% (6/745) of patients, whereas in novel candidate genes, these variants accounted for ~ 0.53% (5/935) and ~ 0.13% (1/745) of patients, respectively. Inherited or state unknown variants in all candidate genes were detected in ~ 35.75% (394/1102) of all patients (Table 3).

Functional Characteristics of Prioritized Candidate Genes

We next employed MetaScape [58] to perform functional enrichment analysis. As expected, these candidate genes were significantly enriched in NDD-associated pathways, such as synapse organization, covalent chromatin modification, head development, behavior, and regulation of ion transport [17, 59–63] (Fig. S3). Interestingly, the novel candidate genes

Table 3 The contribution of prioritized candidate genes to our Chinese probands

Class	Known candidate genes (<i>n</i> = 195)	Novel candidate genes (<i>n</i> = 17)
$FDR_{TADA_combined} < 0.05$ (<i>n</i> = 193)	180	13 genes: <i>MSL2</i> , <i>SPAG9</i> , <i>POLR3A</i> , <i>CYP27C1</i> , <i>DCX</i> , <i>GALNT18</i> , <i>ITSN1</i> , <i>LRR4</i> , <i>PSD3</i> , <i>SMAD6</i> , <i>SPG7</i> , <i>ST3GAL6</i> , <i>UBR3</i>
$0.05 \leq FDR_{TADA_combined} < 0.1$ (<i>n</i> = 15)	12	3 genes: <i>DNAH17</i> , <i>RRAGC</i> , <i>YTHDC1</i>
Genes with X-linked hemizygous (<i>n</i> = 6)	5	1 gene: <i>PLXNA3</i>
Chinese probands with DNVs (<i>n</i> = 935)	94 (10.05%)	5 (0.53%) Subtotal: 99 (10.59%)
Chinese probands with X-linked hemizygous variants (<i>n</i> = 745)	6 (0.81%)	1 (0.13%) Subtotal: 7 (0.94%)
Chinese probands with inherited or state unknown variants (<i>n</i> = 1102)	345 (31.31%)	49 (4.45%) Subtotal: 394 (35.75%)

PTCHD1 and *ARHGEF9* were in two groups, including $FDR_{TADA_combined} < 0.05$ and genes with X-linked hemizygous

were also involved in similar biological pathways. For example, *MSL2* was implicated in covalent chromatin modification, *LRRC4* was involved in chemical synaptic transmission, and *PLXNA3* was associated with head development and axonogenesis.

We also performed functional cell-specific enrichment analyses [64, 65] to investigate whether candidate genes were associated with specific tissues or cells. By analyzing the mouse transcriptomic profiling datasets of different developmental stages and brain regions, we found that the 212 NDD candidate genes tended to be enriched in the cortex and striatum during the middle fetal stage (Fig. S4a). In the cell-specific enrichment analyses, we observed a highly significant enrichment in *Drd1* + and *Drd2* + spiny neurons of neostriatum and rods (Fig. S4b), similar to data reported in a previous study [21]. These results suggest that the 212 NDD candidate genes are functionally associated with the etiology of NDDs.

Functional Network Analysis Between Known and Novel Candidate Genes

To investigate correlations between novel and known candidate genes, we performed a permutation test to estimate the relationship between these genes based on co-expression gene pairs identified from the BrainSpan atlas. We found that 15 of the 17 novel candidate genes ($p = 2.35 \times 10^{-3}$, permutation test, Fig. S5a) were co-expressed with 285 known candidate genes ($p = 3.72 \times 10^{-4}$, permutation test, Fig. S5b), with 523 connections between them ($p = 8.80 \times 10^{-5}$, permutation test, Fig. S5c), suggesting that the novel candidate genes are significantly co-expressed with the known candidate genes and are more likely to be related to the pathology of NDDs.

To further investigate the functional relationship between novel and known candidate genes, we constructed a functional network by integrating PPI data from IntAct and brain expression data from BrainSpan. Only known candidate genes directly interacting/co-expressed with at least two novel candidate genes were added to the network. The co-expressed/PPI network encompassed 159 genes, including 11 novel candidate genes and 148 known candidate genes (Fig. 2). These genes were enriched in several biological processes known to be related to NDDs, such as covalent chromatin modification (GO:0016569, $p = 2.35 \times 10^{-12}$, Fisher's exact test), chemical synaptic transmission (GO:00072686, $p = 7.31 \times 10^{-12}$, Fisher's exact test), and brain development (GO:0007420, $p = 1.23 \times 10^{-7}$, Fisher's exact test). Furthermore, these genes showed a significant enrichment of previously reported gene sets: FMRP targets [66] ($p < 2.22 \times 10^{-16}$, Fisher's exact test), and genes essential in mice [67] ($p < 2.22 \times 10^{-16}$, Fisher's exact test). It is worth noting that six novel candidate genes (*UBR3*, *PLXNA3*, *ITSN1*, *MSL2*, *LRRC4*, and *SPAG9*) were significantly involved in functional

clusters known to be associated with NDDs. For example, *MSL2*, *UBR3*, *SPAG9*, and *PLXNA3* are involved in chromatin organization, while *ITSN1*, *SPAG9*, and *PLXNA3* have been implicated in nervous system development. In addition, we found that nine novel candidate genes showed co-expression/interaction with more than two known candidate genes (Fig. S6). For example, *POLR3A*, which was the most frequently connected novel gene, was co-expressed/interacted with 99 known candidate genes. Other genes (*LRRC4*, *ITSN1*, *UBR3*, *PLXNA3*, *MSL2*, *SPAG9*, *PSD3*, and *YTHDC1*) were connected with more than five candidate genes. These results suggest that novel candidate genes are functionally associated with known risk genes and that these 11 novel candidate genes may have a stronger influence in the etiology of NDDs.

Discussion

Previous studies have provided evidences that targeted sequencing and integrative analyses play a critical role in the discovery of novel candidate genes [21–24]. Based on targeted sequencing of 547 target genes, we identified 108 DNVs, accounting for ~ 10.91% (102/935) of our Chinese cohort. Among the genes with DNVs in our study, most genes have been reported to be associated with NDDs in previous studies, suggesting that genes identified in our Chinese cohort are relevant in other populations. Among the 108 DNVs, 60 DNVs were not recorded in public databases (including dbSNP, gnomAD, Gene4Denovo [39], and PubMed), consistent with the findings of Georgi et al., who reported that recurrently mutated amino acid sites in genes are rarely detected [68]. These results highlight the role of DNVs in the genetic heterogeneity of NDDs. In addition, we identified 21 genes with multiple DNVs present in 51 Chinese patients, and our analyses suggest that these genes display significant intolerance to damaging variants. Consistent with previous ASD study [19], *SCN2A* is the most frequent gene with multiple DNVs in the Chinese population. Compared with the known candidate genes, 20 of 21 genes with multiple DNVs were classified as known candidate genes. Notably, this study is the first to report *TCF20* (with its three de novo PTVs) as a candidate NDD gene in a Chinese cohort.

By integrating information regarding DNVs from public datasets with our results, we prioritized 212 candidate genes, confirming that combing public NDD datasets is beneficial for the discovery of candidate genes [22–24]. Consistent with our analyses of genes with multiple DNVs, the 212 identified candidate genes were more intolerant of damaging variants. In addition, we demonstrated that the 212 candidate genes are closely associated with the etiology of NDDs from the perspective of biological pathway and functional cell-specific enrichment analyses. These results suggest that most of the




Fig. 2 Functional network of novel and known candidate genes. Based on co-expression data from BrainSpan and PPI data from IntAct, 159 candidate genes formed a large interconnected functional network, mainly involving the following major functional clusters: chromatin organization, essential genes, nervous system development, FMRP targets,

behavior, and synapse organization. Novel candidate genes are in red circles and known candidate genes are in green circles. Different line types between nodes represent the interactions existing in BrainSpan or IntAct or in both BrainSpan and IntAct

212 candidate genes identified in our study truly contribute to NDDs and they are worth validating in genetic functional studies or replicating in cohorts.

In this study, we prioritized 17 novel candidate genes and revealed similar functional characteristics between these novel candidate genes and other known candidate genes. Through functional network analysis, we observed that novel candidate genes frequently interacted/were co-expressed with known candidate genes, and genes in the network were enriched in NDD-associated clusters, as described in previous studies [4, 17, 23, 69, 70]. Interestingly, six novel candidate genes were closely connected with known candidate genes and were involved in NDD-associated clusters, suggesting that these novel candidate genes are more likely to be associated with NDDs. For example, *ITSN1* (with an FDR value < 0.01 in the TADA analysis) was involved in nervous system development and synapse organization and connected with 64 known candidate genes. Jakob et al. reported that loss of the signaling scaffold intersectin 1 (*ITSN1*) in mice led to defective neuronal migration and ablates Reelin stimulation of hippocampal long-term potentiation [71]. Our analyses revealed that *SPAG9* carried a de novo PTV in our cohort and it is reportedly overexpressed in human astrocytoma which arises from neural progenitor cells in the central nervous system [72]. We also found that *SPAG9* was implicated in chromatin organization and co-expressed/interacted with eight known candidate genes, further supporting a role for this gene in NDDs. Among the novel candidate genes, *MSL2* was a

particularly interesting gene that carried two de novo PTVs in our Chinese cohort. Except for the multiple DNVs in *MSL2*, it was the gene that frequently co-expressed/interacted with known candidate genes and was involved in the functional cluster of chromatin organization. Iossifov et al. reported a de novo *MSL2* PTV in a patient with autism [69], further suggesting that *MSL2* may be strongly linked to NDDs. We expect to have a large cohort study or functional experiments to validate this possibility in future studies.

Despite our best efforts to comprehensively integrate public data with our cohort data to discover novel risk genes, our study was still bound by some limitations. First, while we discovered that novel candidate genes identified in our study are functionally associated with known candidate genes at multiple levels, these novel candidate genes lack strong evidences that support their roles in NDDs. We expect that these novel candidate genes could be validated in larger cohorts or through functional genetic experiments. Second, the sample size of our Chinese cohort is not large enough, and future studies need to recruit more volunteers to improve the statistical power and allow for comparisons with variant patterns of other populations. Third, since NDDs are characterized by high clinical and genetic heterogeneity, and multiple risk factors contribute to the etiology of these disorders, candidate gene analyses can only partially elucidate the processes underlying NDDs. Further studies are needed to integrate the impact of distinct influences such as epigenetic, environmental, and genetic factors.

In summary, our study demonstrates that integrating DNVs from multiple NDD-related studies can help in the identification of risk genes. We highlight that the pattern of DNVs in Chinese cohorts is relevant to other populations. Furthermore, we provide evidence at multiple levels that novel candidate genes are functionally associated with known candidate genes. Finally, our study describes new high-confidence risk genes that should aid the study of the NDD etiology and we expect that these genes will be worth analyzing in large cohorts or being validated in genetic functional experiments.

Abbreviations ASD, Autism spectrum disorder; Dmis, Deleterious missense variant; DNV, De novo variant; FDR, False discovery rate; NDD, Neurodevelopmental disorder; PPI, Protein-protein interaction; PTV, Protein-truncating variant; RIV, Rare inherited variant; WES, Whole-exome sequencing; WGS, Whole-genome sequencing

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s12035-021-02377-y>.

Availability of Data and Materials The datasets supporting the results and conclusions of this manuscript are included within the article and its Supplementary Material. All other datasets used and analyzed during the current study are available from the corresponding authors on reasonable request.

Funding This work was supported by the Key Research and Development Program of Guangdong Province (No. 2018B030335001), the National Natural Science Foundation of China (81730036, 81525007, and 81801133), Young Elite Scientist Sponsorship Program by CAST (YESS) (2018QNRC001), Guangzhou Key Project in “Early diagnosis and treatment of autism spectrum disorders” (202007030002), Science and Technology Major Project of Hunan Provincial Science and Technology Department (2018SK1030), Natural Science Foundation of Hunan Province for outstanding Young Scholars (2020JJ3059), and Innovation-Driven Project of Central South University (20180033040004).

Declarations

Ethics Approval This study was approved by the Institutional Review Board of the State Key Laboratory of Medical Genetics, School of Life Sciences, Central South University, Changsha, Hunan, China. All subjects who participated in this study provided informed consent prior to sample collection.

Consent to Participate Not applicable

Consent for Publication Not applicable

Conflict of Interest The authors declare no competing interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included

in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Thapar A, Cooper M, Rutter M (2017) Neurodevelopmental disorders. *Lancet Psychiatry* 4(4):339–346. [https://doi.org/10.1016/S2215-0366\(16\)30376-5](https://doi.org/10.1016/S2215-0366(16)30376-5)
2. Mullin AP, Gokhale A, Moreno-De-Luca A et al (2013) Neurodevelopmental disorders: mechanisms and boundary definitions from genomes, interactomes and proteomes. *Transl Psychiatry* 3:e329. <https://doi.org/10.1038/tp.2013.108>
3. Tarlungeanu DC, Novarino G (2018) Genomics in neurodevelopmental disorders: an avenue to personalized medicine. *Exp Mol Med* 50(8):100–107. <https://doi.org/10.1038/s12276-018-0129-7>
4. Stessman HAF, Xiong B, Coe BP, Wang T, Hoekzema K, Fencikova M, Kvarnung M, Gerds J et al (2017) Targeted sequencing identifies 91 neurodevelopmental-disorder risk genes with autism and developmental-disability biases. *Nat Genet* 49(4):515–526. <https://doi.org/10.1038/ng.3792>
5. Wilfert AB, Sulovari A, Turner TN, Coe BP, Eichler EE (2017) Recurrent de novo mutations in neurodevelopmental disorders: properties and clinical implications. *Genome Med* 9(1):101. <https://doi.org/10.1186/s13073-017-0498-x>
6. Gilissen C, Hehir-Kwa JY, Thung DT, van de Vorst M, van Bon BWM, Willemsen MH, Kwint M, Janssen IM et al (2014) Genome sequencing identifies major causes of severe intellectual disability. *Nature* 511(7509):344–347. <https://doi.org/10.1038/nature13394>
7. Sandin S, Lichtenstein P, Kuja-Halkola R, Hultman C, Larsson H, Reichenberg A (2017) The heritability of autism spectrum disorder. *JAMA* 318(12):1182–1184. <https://doi.org/10.1001/jama.2017.12141>
8. Tick B, Bolton P, Happe F et al (2016) Heritability of autism spectrum disorders: a meta-analysis of twin studies. *J Child Psychol Psychiatry* 57(5):585–595. <https://doi.org/10.1111/jcpp.12499>
9. Sullivan PF, Kendler KS, Neale MC (2003) Schizophrenia as a complex trait: evidence from a meta-analysis of twin studies. *Arch Gen Psychiatry* 60(12):1187–1192. <https://doi.org/10.1001/archpsyc.60.12.1187>
10. Chen T, Giri M, Xia Z, Subedi YN, Li Y (2017) Genetic and epigenetic mechanisms of epilepsy: a review. *Neuropsychiatr Dis Treat* 13:1841–1859. <https://doi.org/10.2147/NDT.S142032>
11. Li Y, Jia X, Wu H, Xun G, Ou J, Zhang Q, Li H, Bai T et al (2018) Genotype and phenotype correlations for SHANK3 de novo mutations in neurodevelopmental disorders. *Am J Med Genet A* 176(12):2668–2676. <https://doi.org/10.1002/ajmg.a.40666>
12. Stessman HAF, Willemsen MH, Fencikova M, Penn O, Hoischen A, Xiong B, Wang T, Hoekzema K et al (2016) Disruption of POGZ is associated with intellectual disability and autism spectrum disorders. *Am J Hum Genet* 98(3):541–552. <https://doi.org/10.1016/j.ajhg.2016.02.004>
13. Bernier R, Golzio C, Xiong B, Stessman HA, Coe BP, Penn O, Witherspoon K, Gerds J et al (2014) Disruptive CHD8 mutations define a subtype of autism early in development. *Cell* 158(2):263–276. <https://doi.org/10.1016/j.cell.2014.06.017>
14. Zhou WZ, Zhang J, Li Z, Lin X, Li J, Wang S, Yang C, Wu Q et al (2019) Targeted resequencing of 358 candidate genes for autism spectrum disorder in a Chinese cohort reveals diagnostic potential

- and genotype-phenotype correlations. *Hum Mutat* 40(6):801–815. <https://doi.org/10.1002/humu.23724>
15. Rees E, Carrera N, Morgan J, Hambridge K, Escott-Price V, Pocklington AJ, Richards AL, Pardiñas AF et al (2019) Targeted sequencing of 10,198 samples confirms abnormalities in neuronal activity and implicates voltage-gated sodium channels in schizophrenia pathogenesis. *Biol Psychiatry* 85(7):554–562. <https://doi.org/10.1016/j.biopsych.2018.08.022>
 16. Fromer M, Pocklington AJ, Kavanagh DH, Williams HJ, Dwyer S, Gormley P, Georgieva L, Rees E et al (2014) De novo mutations in schizophrenia implicate synaptic networks. *Nature* 506(7487):179–184. <https://doi.org/10.1038/nature12929>
 17. De Rubeis S, He X, Goldberg AP et al (2014) Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature* 515(7526):209–215. <https://doi.org/10.1038/nature13772>
 18. Mirzaa GM, Chong JX, Piton A et al (2020) De novo and inherited variants in ZNF292 underlie a neurodevelopmental disorder with features of autism spectrum disorder. *Genet Med* 22(3):538–546. <https://doi.org/10.1038/s41436-019-0693-9>
 19. Wang T, Guo H, Xiong B, Stessman HAF, Wu H, Coe BP, Turner TN, Liu Y et al (2016) De novo genic mutations among a Chinese autism spectrum disorder cohort. *Nat Commun* 7:13316–13316. <https://doi.org/10.1038/ncomms13316>
 20. Guo H, Wang T, Wu H, Long M, Coe BP, Li H, Xun G, Ou J et al (2018) Inherited and multiple de novo mutations in autism/developmental delay risk genes suggest a multifactorial model. *Molecular autism* 9:64–64. <https://doi.org/10.1186/s13229-018-0247-z>
 21. Coe BP, Stessman HAF, Sulovari A, Geisheker MR, Bakken TE, Lake AM, Dougherty JD, Lein ES et al (2019) Neurodevelopmental disease genes implicated by de novo mutation and copy number variation morbidity. *Nat Genet* 51(1):106–116. <https://doi.org/10.1038/s41588-018-0288-4>
 22. Nguyen HT, Bryois J, Kim A, Dobbyn A, Huckins LM, Munoz-Manchado AB, Ruderfer DM, Genovese G et al (2017) Integrated Bayesian analysis of rare exonic variants to identify risk genes for schizophrenia and neurodevelopmental disorders. *Genome Med* 9(1):114. <https://doi.org/10.1186/s13073-017-0497-y>
 23. Takata A, Miyake N, Tsurusaki Y, Fukai R, Miyatake S, Koshimizu E, Kushima I, Okada T et al (2018) Integrative analyses of de novo mutations provide deeper biological insights into autism spectrum disorder. *Cell Rep* 22(3):734–747. <https://doi.org/10.1016/j.celrep.2017.12.074>
 24. Gonzalez-Mantilla AJ, Moreno-De-Luca A, Ledbetter DH et al (2016) A cross-disorder method to identify novel candidate genes for developmental brain disorders. *JAMA Psychiatry* 73(3):275–283. <https://doi.org/10.1001/jamapsychiatry.2015.2692>
 25. Li J, Wang L, Guo H, Shi L, Zhang K, Tang M, Hu S, Dong S et al (2017) Targeted sequencing and functional analysis reveal brain-size-related genes and their networks in autism spectrum disorders. *Mol Psychiatry* 22(9):1282–1290. <https://doi.org/10.1038/mp.2017.140>
 26. Li J, Wang L, Yu P et al (2017) Vitamin D-related genes are subjected to significant de novo mutation burdens in autism spectrum disorder. *Am J Med Genet B Neuropsychiatr Genet*:174
 27. Zhang Y, Li N, Li C, Zhang Z, Teng H, Wang Y, Zhao T, Shi L et al (2020) Genetic evidence of gender difference in autism spectrum disorder supports the female-protective effect. *Transl Psychiatry* 10(1):4. <https://doi.org/10.1038/s41398-020-0699-8>
 28. Li J, Cai T, Jiang Y, Chen H, He X, Chen C, Li X, Shao Q et al (2016) Genes with de novo mutations are shared by four neuropsychiatric disorders discovered from NPdenovo database. *Mol Psychiatry* 21(2):290–297. <https://doi.org/10.1038/mp.2015.40>
 29. Zhao G, Li K, Li B, Wang Z, Fang Z, Wang X, Zhang Y, Luo T et al (2019) Gene4Denovo: an integrated database and analytic platform for de novo mutations in humans. *Nucleic Acids Res* 48(D1):D913–D926. <https://doi.org/10.1093/nar/gkz923>
 30. Abrahams BS, Arking DE, Campbell DB et al (2013) SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). *Mol Autism* 4(1):36. <https://doi.org/10.1186/2040-2392-4-36>
 31. Li J, Hu S, Zhang K, Shi L, Zhang Y, Zhao T, Wang L, He X et al (2018) A comparative study of the genetic components of three subcategories of autism spectrum disorder. *Mol Psychiatry* 24:1720–1731. <https://doi.org/10.1038/s41380-018-0081-x>
 32. Wu J, Yu P, Jin X, Xu X, Li J, Li Z, Wang M, Wang T et al (2018) Genomic landscapes of Chinese sporadic autism spectrum disorders revealed by whole-genome sequencing. *J Genet Genomics* 45(10):527–538. <https://doi.org/10.1016/j.jgg.2018.09.002>
 33. Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetjournal*, vol 17, No 1. Next Generation Sequencing Data Analysis. <https://doi.org/10.14806/ej.17.1.200>
 34. Li H, Durbin R (2010) Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26(5):589–595. <https://doi.org/10.1093/bioinformatics/btp698>
 35. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G et al (2009) The sequence alignment/map format and SAMtools. *Bioinformatics* 25(16):2078–2079. <https://doi.org/10.1093/bioinformatics/btp352>
 36. Wang K, Li M, Hakonarson H (2010) ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* 38(16):e164. <https://doi.org/10.1093/nar/gkq603>
 37. Li J, Zhao T, Zhang Y, Zhang K, Shi L, Chen Y, Wang X, Sun Z (2018) Performance evaluation of pathogenicity-computation methods for missense variants. *Nucleic Acids Res* 46(15):7793–7804. <https://doi.org/10.1093/nar/gky678>
 38. He X, Sanders SJ, Liu L, de Rubeis S, Lim ET, Sutcliffe JS, Schellenberg GD, Gibbs RA et al (2013) Integrated model of de novo and inherited genetic variants yields greater power to identify risk genes. *PLoS Genet* 9(8):e1003671. <https://doi.org/10.1371/journal.pgen.1003671>
 39. Zhao G, Li K, Li B, Wang Z, Fang Z, Wang X, Zhang Y, Luo T et al (2019) Gene4Denovo: an integrated database and analytic platform for de novo mutations in humans. *Nucleic Acids Res*. <https://doi.org/10.1093/nar/gkz923>
 40. Sanders SJ, He X, Willsey AJ, Ercan-Sencicek AG, Samocha KE, Cicek AE, Murtha MT, Bal VH et al (2015) Insights into autism spectrum disorder genomic architecture and biology from 71 risk loci. *Neuron* 87(6):1215–1233. <https://doi.org/10.1016/j.neuron.2015.09.016>
 41. Lelieveld SH, Reijnders MR, Pfündt R et al (2016) Meta-analysis of 2,104 trios provides support for 10 new genes for intellectual disability. *Nat Neurosci* 19(9):1194–1196. <https://doi.org/10.1038/nn.4352>
 42. McRae JF, Clayton S, Fitzgerald TW et al (2017) Prevalence and architecture of de novo mutations in developmental disorders. *Nature* 542(7642):433–438. <https://doi.org/10.1038/nature21062>
 43. Stessman HA, Xiong B, Coe BP et al (2017) Targeted sequencing identifies 91 neurodevelopmental-disorder risk genes with autism and developmental-disability biases. *Nat Genet* 49(4):515–526. <https://doi.org/10.1038/ng.3792>
 44. RK CY, Merico D, Bookman M et al (2017) Whole genome sequencing resource identifies 18 new candidate genes for autism spectrum disorder. *Nat Neurosci* 20(4):602–611. <https://doi.org/10.1038/nn.4524>
 45. Satterstrom FK, Kosmicki JA, Wang J, Breen MS, de Rubeis S, An JY, Peng M, Collins R et al (2020) Large-scale exome sequencing study implicates both developmental and functional changes in the

- neurobiology of autism. *Cell* 180(3):568–584 e23. <https://doi.org/10.1016/j.cell.2019.12.036>
46. Lek M, Karczewski KJ, Minikel EV et al (2016) Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 536(7616):285–291. <https://doi.org/10.1038/nature19057>
 47. Petrovski S, Wang Q, Heinzen EL, Allen AS, Goldstein DB (2013) Genic intolerance to functional variation and the interpretation of personal genomes. *PLoS Genet* 9(8):e1003709. <https://doi.org/10.1371/journal.pgen.1003709>
 48. Nava C, Lamari F, Héron D, Mignot C, Rastetter A, Keren B, Cohen D, Faudet A et al (2012) Analysis of the chromosome X exome in patients with autism spectrum disorders identified novel candidate genes, including TMLHE. *Transl Psychiatry* 2(10):e179–e179. <https://doi.org/10.1038/tp.2012.102>
 49. Lim ET, Raychaudhuri S, Sanders SJ et al (2013) Rare complete knockouts in humans: population distribution and significant role in autism spectrum disorders. *Neuron* 77(2):235–242. <https://doi.org/10.1016/j.neuron.2012.12.029>
 50. Toma C, Torricco B, Hervas A et al (2014) Exome sequencing in multiplex autism families suggests a major role for heterozygous truncating mutations. *Mol Psychiatry* 19(7):784–790. <https://doi.org/10.1038/mp.2013.106>
 51. Al-Mubarak B, Abouelhoda M, Omar A et al (2017) Whole exome sequencing reveals inherited and de novo variants in autism spectrum disorder: a trio study from Saudi families. *Sci Rep* 7(1):5679. <https://doi.org/10.1038/s41598-017-06033-1>
 52. Jin SC, Homsy J, Zaidi S, Lu Q, Morton S, DePalma SR, Zeng X, Qi H et al (2017) Contribution of rare inherited and de novo variants in 2,871 congenital heart disease probands. *Nat Genet* 49(11):1593–1601. <https://doi.org/10.1038/ng.3970>
 53. Sullivan PF, Geschwind DH (2019) Defining the genetic, genomic, cellular, and diagnostic architectures of psychiatric disorders. *Cell* 177(1):162–183. <https://doi.org/10.1016/j.cell.2019.01.015>
 54. Krumm N, Turner TN, Baker C et al Excess of rare, inherited truncating mutations in autism. *Nat Genet* 47(6):582–588
 55. Ben-David E, Shifman S (2013) Combined analysis of exome sequencing points toward a major role for transcription regulation during brain development in autism. *Mol Psychiatry* 18(10):1054–1056. <https://doi.org/10.1038/mp.2012.148>
 56. Hoischen A, Krumm N, Eichler EE (2014) Prioritization of neurodevelopmental disease genes by discovery of new mutations. *Nat Neurosci* 17(6):764–772. <https://doi.org/10.1038/nn.3703>
 57. Liu L, Sabo A, Neale BM, Nagaswamy U, Stevens C, Lim E, Bodea CA, Muzny D et al (2013) Analysis of rare, exonic variation amongst subjects with autism spectrum disorders and population controls. *PLoS Genet* 9(4):e1003443. <https://doi.org/10.1371/journal.pgen.1003443>
 58. Tripathi S, Pohl MO, Zhou Y, Rodriguez-Frandsen A, Wang G, Stein DA, Moulton HM, DeJesus P et al (2015) Meta- and orthogonal integration of influenza “OMICs” data defines a role for UBR4 in virus budding. *Cell Host Microbe* 18(6):723–735. <https://doi.org/10.1016/j.chom.2015.11.002>
 59. Bourgeron T (2015) From the genetic architecture to synaptic plasticity in autism spectrum disorder. *Nat Rev Neurosci* 16(9):551–563. <https://doi.org/10.1038/nrn3992>
 60. Mullins C, Fishell G, Tsien RW (2016) Unifying views of autism spectrum disorders: a consideration of autoregulatory feedback loops. *Neuron* 89(6):1131–1156. <https://doi.org/10.1016/j.neuron.2016.02.017>
 61. de la Torre-Ubieta L, Won H, Stein JL, Geschwind DH (2016) Advancing the understanding of autism disease mechanisms through genetics. *Nat Med* 22(4):345–361. <https://doi.org/10.1038/nm.4071>
 62. Ebert DH, Greenberg ME (2013) Activity-dependent neuronal signalling and autism spectrum disorder. *Nature* 493(7432):327–337. <https://doi.org/10.1038/nature11860>
 63. Nanou E, Catterall WA (2018) Calcium channels, synaptic plasticity, and neuropsychiatric disease. *Neuron* 98(3):466–481. <https://doi.org/10.1016/j.neuron.2018.03.017>
 64. Xu X, Wells AB, O'Brien DR, Nehorai A, Dougherty JD (2014) Cell type-specific expression analysis to identify putative cellular mechanisms for neurogenetic disorders. *J Neurosci* 34(4):1420–1431. <https://doi.org/10.1523/JNEUROSCI.4488-13.2014>
 65. Dougherty JD, Schmidt EF, Nakajima M, Heintz N (2010) Analytical approaches to RNA profiling data for the identification of genes enriched in specific cells. *Nucleic Acids Res* 38(13):4218–4230. <https://doi.org/10.1093/nar/gkq130>
 66. Darnell JC, Van Driesche SJ, Zhang C et al (2011) FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* 146(2):247–261. <https://doi.org/10.1016/j.cell.2011.06.013>
 67. Georgi B, Voight BF, Bucan M (2013) From mouse to human: evolutionary genomics analysis of human orthologs of essential genes. *PLoS Genet* 9(5):e1003484. <https://doi.org/10.1371/journal.pgen.1003484>
 68. Geisheker MR, Heymann G, Wang T, Coe BP, Turner TN, Stessman HAF, Hoekzema K, Kvarnung M et al (2017) Hotspots of missense mutation identify neurodevelopmental disorder genes and functional domains. *Nat Neurosci* 20(8):1043–1051. <https://doi.org/10.1038/nn.4589>
 69. Iossifov I, O’Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D, Stessman HA, Witherspoon KT et al (2014) The contribution of de novo coding mutations to autism spectrum disorder. *Nature* 515(7526):216–221. <https://doi.org/10.1038/nature13908>
 70. Zoghbi HY, Bear MF (2012) Synaptic dysfunction in neurodevelopmental disorders associated with autism and intellectual disabilities. *Cold Spring Harb Perspect Biol* 4(3). <https://doi.org/10.1101/cshperspect.a009886>
 71. Jakob B, Kochlamazashvili G, Japel M et al (2017) Intersectin 1 is a component of the Reelin pathway to regulate neuronal migration and synaptic plasticity in the hippocampus. *Proc Natl Acad Sci U S A* 114(21):5533–5538. <https://doi.org/10.1073/pnas.1704447114>
 72. Yi F, Ni W, Liu W, Pan X, Han X, Yang L, Kong X, Ma R et al (2013) SPAG9 is overexpressed in human astrocytoma and promotes cell proliferation and invasion. *Tumour Biol* 34(5):2849–2855. <https://doi.org/10.1007/s13277-013-0845-5>